

Public awareness, attitudes and understanding towards epilepsy in Kelantan, Malaysia

Vimalan RAMASUNDRUM, *Zabidi Azhar MOHD HUSSIN, Chong Tin TAN

*Department of Medicine, University of Malaya, Kuala Lumpur, *Department of Pediatrics, Universiti Sains Malaysia, Kelantan*

Abstract

This is a survey on the public awareness, attitudes and knowledge toward epilepsy in Kota Bharu, Kelantan, Malaysia. Kelantan is a rural state with predominantly Malay population. The survey was based on a questionnaire conducted as one-to-one interview by medical students in the public places. There were 839 respondents, the mean age was 27 years, 65% were married. Sixty-eight percent had received secondary education or above. The respondents have high awareness of epilepsy, with 91% having heard or knew about epilepsy and 56% knew an epileptic person. The attitude towards epilepsy is more negative than similar studies done earlier among Malaysian Chinese. Twenty percent objected to their children associating with a person who sometimes had seizures, 48% objected to their children marrying someone who sometimes had seizures, and 58% thought that people with epilepsy should not be employed in jobs like other people. Only 30% identified epilepsy as a brain disease or disorder, and 69% identified hereditary as the cause. The respondents were not familiar with nonconvulsive form of epilepsy with only 18% identified transient change of behavior, and 16% identified loss of memory as manifestations of epilepsy. As for treatment, 52% advocated prayer, 44% asked for a alternative medicine practitioner, and 9% thought that epilepsy need not be treated or was untreatable. In *conclusion*, the survey of a predominantly Malay population in Malaysia showed that the respondents were familiar with epilepsy, but many maintained a negative attitude and had poor knowledge on causation and treatment of epilepsy.

Key Words: Epilepsy, awareness, attitudes, Kelantan, Malaysia

INTRODUCTION

The public awareness and understanding of a medical illness is crucial in the prophylaxis, early treatment, and compliance of the modern therapy in a community. In epilepsy, the lack of it probably contributes to the high treatment gap seen in many developing countries.^{1,2} The related phenomenon is that the patients may exhaust their limited resources in non-evidence based alternative therapies. The public awareness and understanding may also be important in the formulation of public health policy and allocation of fund. An accurate knowledge of the public attitude to epilepsy is also important as misconception and social misunderstanding may affect the quality of life of the patients more than the seizures itself.

In the recent years, surveys to assess the awareness, understanding and epilepsy have been done among the public in some of the Asian countries, including China³, Taiwan⁴, India^{5,6}, Malaysia⁷, Singapore⁸, among the school teachers

in Thailand⁹ and patients in Pakistan.¹⁰ These studies generally showed a similar level of awareness, but more negative attitudes towards epilepsy when compared to the developing countries in the West. The previous Malaysian study was done among ethnic Chinese in an urban and rural area around Kuala Lumpur. As Malaysia is a multi-racial country where ethnic Chinese constitutes 25% of the population, it is important to extend the study to other population groups. This is the report of a survey done in Kota Bharu, Kelantan, in Peninsular Malaysia whose residents are predominantly Malays.

METHODS

The study was performed in Kota Bharu, the capital of Kelantan state. The study consisted of a survey using a standard questionnaire of ten questions. To facilitate comparison, the questionnaire was similar to that used in the Gallup Poll¹¹ and the previous studies in China³, Taiwan⁴, Singapore⁸ and Malaysia.⁷ The

questionnaire was translated into Malay. The survey was conducted as a one-to-one interview by medical students after a period of training. The interviewers were encouraged not to lead or coax the respondents in their answers and only explain the questions when it was necessary to clarify the points. The respondents were randomly selected at public places such as shopping complexes, hospitals and town halls. Subjects who were 13 years or above willing to participate in the questionnaire were recruited.

The data was analyzed using the SPSS version 6.0 with descriptive statistics and the level of significance was set at $p < 0.05$.

RESULTS

Eight hundred and thirty nine subjects participated in the survey. The mean age was 27 years. The male to female sex ratio was 1 : 1.6. The predominant languages spoken were: Malay (81%), Chinese (6.0%), Tamil (0.6%), English (4.3%) and Others (8.0%). Most of the subjects were married (65%), the remaining were single (33%) and divorced or separated (2%). For the married subjects, the number of children was: none (12.5%), 1-2 children (24%), 3-5 children (42%), 6-10 children (19%) and 11-12 children (1.7%). The distribution of the occupation was: secondary students (34%), traders (14%), teachers (6%), government servants (6%), medical students (4.5%), farmers (1.5%), fishermen (1.5%), factory workers (1.5%) and others (30%). The level of education was: secondary education (68%), university education (12%), primary education (6%), and no formal education (14%). The residence of the respondents was: in or around the town (74%), rural areas (16%) and unknown (9%).

Familiarity with epilepsy

The responses to the questions related to familiarity with epilepsy are listed in Table 1. Ninety-one percent had heard or knew about epilepsy, 56% knew an epileptic person and 33% had witnessed an attack of epilepsy. Significantly greater number of respondents younger than 50 years and those with secondary or tertiary education have heard or read of epilepsy. Significantly greater number of those older than 50 years, the married and respondents with greater than three children have known someone with epilepsy. Significantly greater number of those older than 50 years and the married have witnessed an epileptic attack.

Attitudes towards epilepsy

The responses to questions related to attitudes towards epilepsy are listed in Table 2. Twenty percent of the respondents objected to the possibility that their children might be playing with an epileptic. The objection was significantly higher among the respondents older than 50 years of age but less among those with tertiary education. Forty-eight percent objected to their children marrying a person who sometimes have seizure. The objection was significantly higher among those with tertiary education and those from the urban area. Fifty-eight percent of the respondents thought that people with epilepsy cannot be employed in jobs like other people. The attitude was significantly more favourable among those from the town area. Twenty-three percent of the respondents believed that epilepsy was a form of insanity.

Understanding of epilepsy

The understanding of epilepsy elucidated was in the area of knowledge of its cause, presentation and treatment. The responses are as listed in Tables 3-5.

DISCUSSION

This study was done in Kota Bharu, the state capital of Kelantan. Kelantan is located at the North Eastern part of Peninsular Malaysia. The state has a population of one and half million. Kota Bahru itself has 400,000 population. Ninety-five percent of the Kelantan population are Malays. Ethnic Chinese, Thais and Indians constitute the rest of the population. Whereas the urban and rural population are equally divided in Malaysia, two third of the population in Kelantan is rural. When compared with the states in the west coast of Peninsular Malaysia, the economy in Kelantan is also less developed. The study subjects in this study is thus representative of the town population from a rural Malay state in Malaysia.

The responses to questions related to familiarity with epilepsy show that the study population has high awareness of epilepsy. The high positive response rate of 91% for having heard or read about epilepsy is similar to the other Asian studies.³⁻⁸ As for the question of whether the respondent has witnessed someone with a seizure, the positive rate was 33%. The corresponding figure in Henan, China was 72%. The disproportionately higher awareness in the Chinese population was attributed to the

TABLE 1: Responses to questions related to familiarity with epilepsy

	No. of response	Q1		Q2		Q3	
		Yes %	No %	Yes %	No %	Yes %	No %
Total	839	91	9	56	44	33	67
Age of respondents (years)							
13-29	565	91	9	52	48	30	70
30-49	181	91	6	61	39	37	63
≥50	93	84	16	68	32	47	53
Sex of respondents							
Male	320	90	10	57	43	33	67
Female	519	92	8	55	45	33	67
Marital status							
Never married	544	92	8	53	47	31	69
Married ^a	295	90	10	61	39	39	61
No. of children							
0	569	92	8	54	46	31	69
1-2	182	88	12	52	48	32	68
≥3	191	91	9	66	34	41	59
Education level							
Primary ^b	76	73	27	64	36	37	63
Secondary	567	91	9	51	49	29	71
Tertiary	196	95	5	67	33	45	55
Residence							
Urban ^c	625	91	9	59	41	33	67
Rural	213	91	9	47	53	32	68
Occupation							
Students	282	91	9	50	50	32	68
Traders, medical students, teachers ^d	252	92	8	72	28	40	60
Farmers ^e	52	93	7	50	50	40	60
Others	253	92	8	48	52	26	74

Three questions were asked. Q1: Have you ever heard of or read about the disease called 'epilepsy' or convulsive seizures (fits)? Q2: Did you ever know anyone who had epilepsy? Q3: Have you ever seen anyone having a seizure?

^a Married include widow, widowers and divorcees

^b Include those with no formal education

^c Urban includes town and perimeter of town

^d Include other government workers

^e Include factory workers

TABLE 2: Responses to questions about attitudes towards epilepsy

	No. of response	Q4		Q5		Q6		Q7	
		Yes %	No %	Yes %	No %	Yes %	No %	Yes %	No %
Total	839	20	80	48	52	42	58	23	77
Age of respondents (years)									
13-29	565	20	80	50	50	42	58	25	75
30-49	181	17	83	45	55	44	56	17	83
≥50	93	30	70	34	66	38	62	21	79
Sex of respondents									
Male	320	21	79	45	55	40	60	16	84
Female	519	20	80	50	50	43	57	26	74
Marital status									
Never married	544	20	80	49	51	40	60	21	79
Married ^a	295	21	79	47	53	43	57	23	77
No. of children									
0	569	19	81	49	51	42	58	24	76
1-2	182	20	80	52	48	38	62	27	73
≥3	56	21	79	43	57	42	58	17	83
Education level									
Primary ^b	76	39	61	37	63	44	56	21	78
Secondary	567	19	81	47	53	40	60	22	77
Tertiary	196	16	84	59	41	48	52	24	76
Residence									
Urban ^c	625	21	79	54	46	44	56	24	76
Rural	214	16	84	31	69	34	66	21	79
Occupation									
Students	282	20	80	55	45	43	57	26	74
Traders, medical students, teachers ^d	252	23	77	53	47	48	52	22	78
Farmers ^e	52	20	80	55	45	40	60	22	78
Others	233	17	83	39	61	35	65	22	78

Four questions were asked. Q4: Would you object to having any of your children in school or at play associate with a person who sometimes had seizures (fits)? Q5: Would you object to having a son or daughter of yours marrying a person who sometimes had seizures? Q6: Do you think people with epilepsy should or should not be employed in jobs like other people? Q7: Do you think epilepsy is a form of insanity or not?

^a Married include widow, widowers and divorcees

^b Include those with no formal education

^c Urban includes town and perimeter of town

^d Include other government workers

^e Include factory workers

Table 3: Responses to the question: “what do you think is the cause of epilepsy?”

Responses	Percentage positive
Hereditary	69
Brain disease or disorder	30
Mental or emotional disorder	24
Don't know	17
Birth defect	13
Blood disorder	6
Unsure of answer	5
Others	2

Table 4: Responses to the question: “What do you think an epileptic attack is?”

Responses	Percentage positive
Shaking, convulsion	81
Loss of consciousness	62
Transient changes of behavior	18
Loss of memory	16
Unsure of answer	7

Table 5: Response to the question: “If your relatives or friends had epilepsy, what kind of treatment would you suggest?”

Responses	Percentage positive
Ask for a medical doctor	78
Ask for God's help	52
Ask for a alternative medicine practitioner	44
“Dan Fang” and other folk medicine	16
Acupuncture	8
No need to treat	7
Don't know what to recommend	5
Ask for a witch doctor	3
Think “epilepsy is untreatable”	2

overpopulated condition and closer interpersonal relationships in Henan.³

As for the questions on attitudes towards epilepsy, the responses were less negative than similar studies in China, but more negative than that of previous studies among Chinese in Malaysia. This is particularly so for those who had lower level of education. Twenty percent would object to their children in school or at play associate with a person with seizure, the corresponding figure was 57% in China³ and 9% among Malaysian Chinese.⁷ Fifty-eight percent of the respondents thought that people with epilepsy should not be employed in jobs like others. The corresponding figure was 53% in China³ but 14% amongst Malaysian Chinese.⁷

Forty-eight percent objected to their children marrying a person who sometimes had seizures. The corresponding figure was 87% in China³ and 43% among the Malaysian Chinese.⁷ The objection to marriage was unusual in that it was more evident among those with tertiary education and the urban residence.

Less than a third were able to attribute epilepsy to a disorder or disease of the brain. This is similar to other Asian studies.^{3,4,7,8} As in the other Asian studies, there was poor awareness of the non-convulsive form of epilepsy with less than a fifth able to identify “transient changes of behavior” and “loss of memory” as manifestations of epileptic seizure.^{3,4,7,8}

The strong emphasis on prayer to supplement

other treatment reflected the highly religious character of the respondents who were predominantly Muslims. In contrast, only 1% of the Malaysian Chinese would invoke the help of God.⁷ Forty-four percent of the respondents would recommend alternative medicine practitioner for treatment. This is again higher than the Malaysian Chinese where 11% recommended herbal medicine doctor.⁷ Nine percent of the respondents thought that epilepsy was untreatable or there was no need to treat epilepsy. Such a negative attitude to treatment of epilepsy is consistent the previous report of children with epilepsy in Kelantan, where 20% did not receive any form of schooling, although primary schooling is free in Malaysia.¹² In a study on epilepsy awareness among school teachers in Thailand, 47% believed that epilepsy was a chronic incurable disease, and 15% preferred to place all children in a special classroom.⁹ Whether the children with epilepsy would continue in school probably depends on the attitude and knowledge of the school teachers as well as the parents. The negative attitude and poor knowledge on epilepsy as shown in this survey is also consistent with the previous report of high treatment gap in the rural part of Malaysia.²

REFERENCES

1. Shorvon SD, Farmer PJ. Epilepsy in developing countries: A review of epidemiological, socio-cultural and treatment aspects. *Epilepsia* 1988;29:S36-54
2. Menon J, Saw A, Tan CT. Epilepsy treatment gap in Sabah and Kuala Lipis, Malaysia. *Neurol J Southeast Asia* 1996;1:75-6.
3. Lai CW, Huang X, Lai YHC, Zhang Z, Liu G, Yang MZ. Survey of public awareness, understanding and attitudes towards epilepsy in Henan Province, China. *Epilepsia* 1990;31:182-7
4. Chung MY, Chang YC, Lai CW. Survey of public awareness, understanding and attitudes towards epilepsy in Taiwan. *Epilepsia* 1995;36:488-93
5. Gambhir SK, Singhi PD, Goel RC. Public awareness, understanding and attitudes toward epilepsy. *Indian J Med Res* 1995;102:34-8.
6. Radhakrishnan K, Pandian JD, Santoshkumar T, Thomas SV, Deetha TD, Sarma PS, Jayachandran D, Mohamed E. Prevalence, knowledge, attitude and practice of epilepsy in Kerala, South India. *Epilepsia* 2000;41:1027-35.
7. Lim KS, Tan LP, Lim KT, Tan CT. Survey of public awareness, understanding and attitudes towards epilepsy in Malaysia. *Neurol J Southeast Asia* 1999;4:31-36
8. Pan APS, Lim SH. Public awareness, attitudes and understanding toward epilepsy among Singaporean Chinese. *Neurol J Southeast Asia* 2000;5:5-10.
9. Kankirawatana P. Epilepsy awareness among schoolteachers in Thailand. *Epilepsia* 1999;40:497-501
10. Aziz H, Akhtr SW, Hasan KZ. Epilepsy in Pakistan: Stigma and psychosocial problems: A population-based epidemiologic study. *Epilepsia* 1997;38:1069-73.
11. Cavenass WP, Gallup GH Jr. A survey on attitudes towards epilepsy in 1979 with an indication of trends over the past thirty years. *Epilepsia* 1980;21:509-18
12. Hassan H, Mohd Hussin ZA. A five year study of childhood epilepsy in Universiti Sains Malaysia Hospital, Kelantan, Malaysia. *Neurol J Southeast Asia* 1996;1:80-1.